



**Description:**

Male impotency is the cause of embracement for many males who suffers silently. Not only physically but they tend to get affected emotionally through the side effects of long term erectile dysfunction. Even now science has admitted that simply treatable misbalance like erectile dysfunction can easily lend someone to the condition of depression. All this occur due to the unacceptability of oneself and even by the partner.

But not anymore [Vidalista 10 Mg Medicine](#) made from main tadalafil can solve the point of erectile dysfunction helping such couple to enjoy their life to the fullest. [vidalista 10 review](#)

Has shown the tremendous response the medication got in various countries and subcontinents. There are numerous success stories.

### **How to take Vidalista 10 mg?**

Take this medicine in the dose and period as advised by your physician. Vidalista 10 Mg works best when taken one and half hour before sexual activity. You will not get an erection until you are sexually stimulated. The erection will go away by itself after a sexual encounter. The effects of the medication can last up to 36 hours, so you can enjoy the effects of the drugs for more than one day. Swallow it as a whole. Do not crush, chew or break it. It may be take with or without food. Seek emergency medical nursing or contact the physician in case of an overdose.

### **How Vidalista 10 Works?**

Erectile dysfunction occurs when your penis doesn't get filled with enough blood during the sexual stimulation. As a result, you can no longer hold your penis for the desired amount of time. But [Vidalista 10 Mg](#) uses the Tadalafil which is a PDE-5 inhibitor. It works by removing the clogged arteries and helps in the relaxation of tissues and muscles.

As a result, the blood vessels can remain open for a longer duration and can supply uniform blood to the penis during sexual intercourse. Thus, you can enjoy pleasurable time. It works like wonder for those suffering this problem, but you should always take it under the right prescription by your doctor.

### **Dosage:**

#### **Missed Dose:**

Take the missed dose as soon as you think of. It can be leapped if it almost time for the next scheduled dose. This applies to shape like Pulmonary Hypertension where the dose regimen is fixed.

#### **Overdose:**

Contact a doctor at once if an overdose is suspected. You might require immediate health check attention if the overdose is considerate.

### **Side Effects:**

- Blurring/Loss of Vision

- Decreased/Loss of Hearing
- Painful / Prolonged Erection
- Breathing Problems
- Diarrhea
- A headache
- Dizziness
- Swelling of Body Parts (face, lips, eyelids etc)
- Body Pain or Sudden Weakness
- Chest Pain
- Skin allergies

**Preacution:**

Don't take [Vidalista 10 Mg Tablets](#) if you are using a nitrate drug for breast pain or heart problems, with nitroglycerin, isosorbide dinitrate, isosorbide mononitrate, and some recreational drugs such as “poppers”. Taking tadalafil with a nitrate medicine can be a reason for sudden and serious decrease in blood pressure.

Some tablets can cause unwanted or hazardous effects when used with tadalafil. Tell your physician about all your existing medicines, especially riociguat (Adempas).

Contact your physician or seek emergency medical attention if your structure is painful or lasts longer than 4 hours. A prolonged erection (priapism) can damage the penis.

Get medical assistance right away if you have nausea, chest pain, or dizziness during sex.

prevent using of tadalafil and get emergency medical assistance if you have sudden eyesight loss.

**Storage:**

Store your [Vidalista 10 Mg Drug](#) at room temperature among 68-77 degrees F (20-25 degrees C) far away from light and humidity. Do not store your medicines in the bathroom. Keep all medicines away from reach of children and pets.

For More Visit Click Here :[medsday.com](http://medsday.com)